



HOLMDEL TOWNSHIP SCHOOL DISTRICT

"A COMMITMENT TO EXCELLENCE"

Office of the Superintendent of Schools
4 Crawford's Corner Road
Holmdel, NJ 07733
tel: 732-946-1800
fax: 732-946-1875

October 30, 2009

Dear Parents/Guardians:

As I am sure you have heard, a new influenza virus, called the 2009 H1N1 influenza virus, was first identified in the United States in late April 2009. The virus has caused illness ranging from mild to severe, including hospitalizations and deaths in adults and children. Many children have gotten 2009 H1N1 infection and there have been large outbreaks in some schools across the country. The CDC has recommended that children and young adults aged six (6) months through twenty-four (24) years be vaccinated against 2009 H1N1 as soon as the vaccine is available. Other groups recommended by the CDC to receive the first doses of 2009 H1N1 influenza vaccine are:


- Pregnant women
- People who live with or care for children younger than 6 months of age
- Health care and emergency medical services workers, and
- People ages twenty-five (25) through sixty-four (64) years who have certain health conditions such as HIV, diabetes, or heart or lung disease.

The Monmouth County Regional Health Commission is working with us to make the 2009 H1N1 influenza vaccine available to Holmdel students. They will hold school-located vaccination clinics on a weekend beginning later this fall. We will let you know the specific dates once we are notified by the Monmouth County Regional Health Commission that the vaccine is on hand. The CDC advises that children under age ten (10) are expected to need two (2) doses of vaccine spaced about roughly four (4) weeks apart. There will be no cost to you for this vaccine.

You can read the Vaccine Information Statements included with this letter about the two (2) types of the vaccine. If you plan to have your child vaccinated, please sign and date the included consent form and return it to the main office at your child's school no later than Friday, November 13, 2009. If, at any time, you change your mind about having your child vaccinated, you can contact your school. Please be advised that giving consent early will allow the Monmouth County Regional Health Commission to estimate the number of vaccinations required for Holmdel's weekend clinic.

If you have any questions about the vaccine or the vaccination clinics, please call your school nurse or the Monmouth County Regional Health Commission at 732-493-9520 from 9 AM to 4:30 PM. Please visit the CDC's 2009 H1N1 influenza web site at <http://www.cdc.gov/h1n1flu/> and also <http://www.cdc.gov/h1n1flu/parents> for more information especially for parents. Your child's health care provider also can answer your questions about the 2009 H1N1 influenza virus and may be able to give your child the seasonal influenza vaccine and the 2009 H1N1 vaccine.

Sincerely,


Barbara Duncan
Superintendent of Schools

Enclosures

For Informational Purposes Only

2009 H1N1 INFLUENZA VACCINE

LIVE, ATTENUATED
(the nasal spray vaccine)

WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.

1 What is 2009 H1N1 influenza?

2009 H1N1 influenza (sometimes called Swine Flu) is caused by a new strain of influenza virus. It has spread to many countries.

Like other flu viruses, 2009 H1N1 spreads from person to person through coughing, sneezing, and sometimes through touching objects contaminated with the virus.

Signs of 2009 H1N1 can include:

- Fatigue
- Fever
- Sore Throat
- Muscle Aches
- Chills
- Coughing
- Sneezing

Some people also have diarrhea and vomiting.

Most people feel better within a week. But some people get pneumonia or other serious illnesses. Some people have to be hospitalized and some die.

2 How is 2009 H1N1 different from regular (seasonal) flu?

Seasonal flu viruses change from year to year, but they are closely related to each other.

People who have had flu infections in the past usually have some immunity to seasonal flu viruses (their bodies have built up some ability to fight off the viruses).

The 2009 H1N1 flu virus is a new virus strain. It is very different from seasonal flu viruses.

Most people have little or no immunity to 2009 H1N1 flu (their bodies are not prepared to fight off the virus).

3 2009 H1N1 influenza vaccine

Vaccines are available to protect against 2009 H1N1 influenza.

- These vaccines are made just like seasonal flu vaccines.
- They are expected to be as safe and effective as seasonal flu vaccines.
- They will not prevent “influenza-like” illnesses caused by other viruses.

- They will not prevent seasonal flu. *You should also get seasonal influenza vaccine, if you want protection from seasonal flu.*

Live, attenuated intranasal vaccine (or LAIV) is sprayed into the nose. **This sheet describes the live, attenuated intranasal vaccine.**

An **inactivated** vaccine is also available, which is given as a shot. It is described in a separate sheet.

The 2009 H1N1 LAIV does not contain thimerosal or other preservatives. It is licensed for people from 2 through 49 years of age.

The vaccine virus is attenuated (weakened) so it will not cause illness.

4 Who should get 2009 H1N1 influenza vaccine and when?

WHO

LAIV is approved for people from 2 through 49 years of age who are not pregnant and do not have certain health conditions (see number 5 below). Groups recommended to receive 2009 H1N1 LAIV first are healthy people who:

- are from 2 through 24 years of age,
- are from 25 through 49 years of age and
 - live with or care for infants younger than 6 months of age, or
 - are health care or emergency medical personnel.

As more vaccine becomes available, other healthy 25 through 49 year olds should also be vaccinated.

Note: While certain groups should not get LAIV – for example pregnant women, people with long-term health problems, and children from 6 months to 2 years of age – it is important that they be vaccinated. They should get the flu shot.

The Federal government is providing this vaccine for receipt on a voluntary basis. However, state law or employers may require vaccination for certain persons.

WHEN

Get vaccinated as soon as the vaccine is available.

Children through 9 years of age should get **two doses** of vaccine, about a month apart. Older children and adults need only one dose.

5 Some people should not get the vaccine or should wait

You should not get 2009 H1N1 LAIV if you have a **severe (life-threatening) allergy** to **eggs**, or to **any other substance in the vaccine**. *Tell the person giving you the vaccine if you have any severe allergies.*

2009 H1N1 LAIV should not be given to the following groups.

- children younger than 2 and adults 50 years and older
- pregnant women,
- anyone with a weakened immune system,
- anyone with a long-term health problem such as
 - heart disease - kidney or liver disease
 - lung disease - metabolic disease such as diabetes
 - asthma - anemia and other blood disorders
- children younger than 5 years with asthma or one or more episodes of wheezing during the past year,
- anyone with certain muscle or nerve disorders (such as cerebral palsy) that can lead to breathing or swallowing problems,
- anyone in close contact with a person with a *severely* weakened immune system (requiring care in a protected environment, such as a bone marrow transplant unit),
- children or adolescents on long-term aspirin treatment.

If you are moderately or severely ill, you might be advised to wait until you recover before getting the vaccine. If you have a mild cold or other illness, there is usually no need to wait.

Tell your doctor if you ever had:

- a life-threatening allergic reaction after a dose of seasonal flu vaccine,
- Guillain-Barré syndrome (a severe paralytic illness also called GBS).

These may not be reasons to avoid the vaccine, but the medical staff can help you decide.

2009 H1N1 LAIV may be given at the same time as most other vaccines. Tell your doctor if you got any other vaccines within the past month or plan to get any within the next month. H1N1 LAIV and seasonal LAIV should not be given together.

6 What are the risks from 2009 H1N1 LAIV?

A vaccine, like any medicine, could cause a serious problem, such as a severe allergic reaction. But the risk of any vaccine causing serious harm, or death, is extremely small.

The risks from 2009 H1N1 LAIV are expected to be similar to those from seasonal LAIV:

Mild problems:

Some children and adolescents 2-17 years of age have reported mild reactions, including:

- runny nose, nasal congestion or cough
- fever
- headache and muscle aches
- wheezing
- abdominal pain or occasional vomiting or diarrhea

Some adults 18-49 years of age have reported:

- runny nose or nasal congestion
- sore throat
- cough, chills, tiredness/weakness
- headache

Severe problems:

- Life-threatening allergic reactions to vaccines are very rare. If they do occur, it is usually within a few minutes to a few hours after the vaccination.
- In 1976, an earlier type of inactivated swine flu vaccine was associated with cases of Guillain-Barré Syndrome (GBS). LAIV has not been linked to GBS.

7 What if there is a severe reaction?

What should I look for?

Any unusual condition, such as a high fever or behavior changes. Signs of a severe allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** the doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your provider to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form. Or you can file this report through the VAERS website at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not provide medical advice.

8 Vaccine injury compensation

If you or your child has a reaction to the vaccine, your ability to sue is limited by law.

However, a federal program has been created to help pay for the medical care and other specific expenses of certain persons who have a serious reaction to this vaccine. For more information about this program, call **1-888-275-4772** or visit the program's website at:

www.hrsa.gov/countermeasurescomp/default.htm.

9 How can I learn more?

- Ask your provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/h1n1flu or www.cdc.gov/flu
 - Visit the web at www.flu.gov



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



2009 H1N1 INFLUENZA VACCINE

INACTIVATED (the “flu shot”)

WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.

1 What is 2009 H1N1 influenza?

2009 H1N1 influenza (also called Swine Flu) is caused by a new strain of influenza virus. It has spread to many countries.

Like other flu viruses, 2009 H1N1 spreads from person to person through coughing, sneezing, and sometimes through touching objects contaminated with the virus.

Signs of 2009 H1N1 can include:

- Fatigue
- Fever
- Sore Throat
- Muscle Aches
- Chills
- Coughing
- Sneezing

Some people also have diarrhea and vomiting.

Most people feel better within a week. But some people get pneumonia or other serious illnesses. Some people have to be hospitalized and some die.

2 How is 2009 H1N1 different from regular (seasonal) flu?

Seasonal flu viruses change from year to year, but they are closely related to each other.

People who have had flu infections in the past usually have some immunity to seasonal flu viruses (their bodies have built up some ability to fight off the viruses).

The 2009 H1N1 flu is a new flu virus. It is very different from seasonal flu viruses.

Most people have little or no immunity to 2009 H1N1 flu (their bodies are not prepared to fight off the virus).

3 2009 H1N1 influenza vaccine

Vaccines are available to protect against 2009 H1N1 influenza.

- These vaccines are made just like seasonal flu vaccines.
- They are expected to be as safe and effective as seasonal flu vaccines.
- They will not prevent “influenza-like” illnesses caused by other viruses.
- They will not prevent seasonal flu. *You should also get seasonal influenza vaccine, if you want to be protected against seasonal flu.*

Inactivated vaccine (vaccine that has killed virus in it) is injected into the muscle, like the annual flu shot. **This sheet describes the inactivated vaccine.**

A **live, intranasal** vaccine (the nasal spray vaccine) is also available. It is described in a separate sheet.

Some inactivated 2009 H1N1 vaccine contains a preservative called thimerosal to keep it free from germs. Some people have suggested that thimerosal might be related to autism. In 2004 a group of experts at the Institute of Medicine reviewed many studies looking into this theory, and found no association between thimerosal and autism. Additional studies since then reached the same conclusion.

4 Who should get 2009 H1N1 influenza vaccine and when?

WHO

Groups recommended to receive 2009 H1N1 vaccine first are:

- Pregnant women
- People who live with or care for infants younger than 6 months of age
- Health care and emergency medical personnel
- Anyone from 6 months through 24 years of age
- Anyone from 25 through 64 years of age with certain chronic medical conditions or a weakened immune system

As more vaccine becomes available, these groups should also be vaccinated:

- Healthy 25 through 64 year olds
- Adults 65 years and older

The Federal government is providing this vaccine for receipt on a voluntary basis. However, state law or employers may require vaccination for certain persons.

WHEN

Get vaccinated as soon as the vaccine is available.

Children through 9 years of age should get **two doses** of vaccine, about a month apart. Older children and adults need only one dose.

5 Some people should not get the vaccine or should wait

You should not get 2009 H1N1 flu vaccine if you have a **severe (life-threatening) allergy to eggs**, or to **any other substance in the vaccine**. *Tell the person giving you the vaccine if you have any severe allergies.*

Also tell them if you have ever had:

- a life-threatening allergic reaction after a dose of seasonal flu vaccine,
- Guillain Barré Syndrome (a severe paralytic illness also called GBS).

These may not be reasons to avoid the vaccine, but the medical staff can help you decide.

If you are moderately or severely ill, you might be advised to wait until you recover before getting the vaccine. If you have a mild cold or other illness, there is usually no need to wait.

Pregnant or breastfeeding women can get inactivated 2009 H1N1 flu vaccine.

Inactivated 2009 H1N1 vaccine may be given at the same time as other vaccines, including seasonal influenza vaccine.

6 What are the risks from 2009 H1N1 influenza vaccine?

A vaccine, like any medicine, could cause a serious problem, such as a severe allergic reaction. But the risk of any vaccine causing serious harm, or death, is extremely small.

The virus in inactivated 2009 H1N1 vaccine has been killed, so you cannot get influenza from the vaccine.

The risks from inactivated 2009 H1N1 vaccine are similar to those from seasonal inactivated flu vaccine:

Mild problems:

- soreness, redness, tenderness, or swelling where the shot was given
- fainting (mainly adolescents)
- headache, muscle aches
- fever
- nausea

If these problems occur, they usually begin soon after the shot and last 1-2 days.

Severe problems:

- Life-threatening allergic reactions to vaccines are very rare. If they do occur, it is usually within a few minutes to a few hours after the shot.
- In 1976, an earlier type of swine flu vaccine was associated with cases of Guillain-Barré Syndrome (GBS). Since then, flu vaccines have not been clearly linked to GBS.

7 What if there is a severe reaction?

What should I look for?

Any unusual condition, such as a high fever or behavior changes. Signs of a severe allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** the doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your provider to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form. Or you can file this report through the VAERS website at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not provide medical advice.

8 Vaccine injury compensation

If you or your child has a reaction to the vaccine, your ability to sue is limited by law.

However, a federal program has been created to help pay for the medical care and other specific expenses of certain persons who have a serious reaction to this vaccine. For more information about this program, call **1-888-275-4772** or visit the program's website at: www.hrsa.gov/countermeasurescomp/default.htm.

9 How can I learn more?

- Ask your provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/h1n1flu or www.cdc.gov/flu
- Visit the web at www.flu.gov



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



**Monmouth County Regional Health Commission #1
2009 H1N1 Influenza Vaccine Consent Form**

Section 1: Information about Child to Receive Vaccine (please print)

STUDENT'S NAME (Last)		(First)	(M.I.)	STUDENT'S DATE OF BIRTH month day year	
PARENT/LEGAL GUARDIAN'S NAME (Last)		(First)	(M.I.)	STUDENT'S AGE	STUDENT'S GENDER M / F
ADDRESS			PARENT/GUARDIAN DAYTIME PHONE NUMBER:		
CITY	STATE	ZIP			
SCHOOL NAME			GRADE		

Section 2: Screening for Vaccine Eligibility

If your child has already been vaccinated with 2009 H1N1 influenza vaccine, please tell us the number of doses and dates of vaccination.

- Dose 1 Date received: month ____ day ____ year ____ Form (please circle): nasal spray shot
- Dose 2 Date received: month ____ day ____ year ____ Form (please circle): nasal spray shot

The following questions will help us to know if your child can get the 2009 H1N1 influenza vaccine. Please mark YES or NO for each question.

A. If you answer "NO" to all four of the following questions, your child can probably get the influenza vaccine. If you answer "YES" to one or more of the following four questions, your child may be able to get the 2009 H1N1 vaccine, but we will contact you to discuss your options.

	YES	NO
1. Does your child have a serious allergy to eggs?	<input type="checkbox"/>	<input type="checkbox"/>
2. Does your child have any other serious allergies? Please list: _____	<input type="checkbox"/>	<input type="checkbox"/>
3. Has your child ever had a serious reaction to a previous dose of flu vaccine?	<input type="checkbox"/>	<input type="checkbox"/>
4. Has your child ever had Guillain-Barré Syndrome (a type of temporary severe muscle weakness) within 6 weeks after receiving a flu vaccine?	<input type="checkbox"/>	<input type="checkbox"/>

B. There are two kinds of 2009 H1N1 influenza vaccine. Your answers to the following questions will help us know which of the two kinds of vaccine your child can get.

	YES	NO
1. Has your child been vaccinated with any vaccine (not just flu) within the past 30 days? Vaccine: _____ Date given: month ____ day ____ year ____	<input type="checkbox"/>	<input type="checkbox"/>
2. Does your child have any of the following: asthma, diabetes (or other type of metabolic disease), or disease of the lungs, heart, kidneys, liver, nerves, or blood?	<input type="checkbox"/>	<input type="checkbox"/>
3. Is your child on long-term aspirin or aspirin-containing therapy (for example, does your child take aspirin every day)?	<input type="checkbox"/>	<input type="checkbox"/>
4. Does your child have a weak immune system (for example, from HIV, cancer, or medications such as steroids or those used to treat cancer)?	<input type="checkbox"/>	<input type="checkbox"/>
5. Is your child pregnant?	<input type="checkbox"/>	<input type="checkbox"/>
6. Does your child have close contact with a person who needs care in a protected environment (for example, someone who has recently had a bone marrow transplant)?	<input type="checkbox"/>	<input type="checkbox"/>

Section 3: Consent

My preference for my (my child's) influenza vaccine is the following:

- Inactivated injectable vaccine only Either injectable influenza vaccine or live attenuated intranasal influenza vaccine
- Live attenuated intranasal influenza vaccine only

CONSENT FOR CHILD'S VACCINATION:

I have read or had explained to me the 2009-2010 Vaccine Information Statement for the 2009 H1N1 influenza vaccine and understand the risks and benefits.

<p>I GIVE CONSENT to Monmouth County Regional Health Commission for my child named at the top of this form to be vaccinated with this vaccine. (If this consent form is not signed, dated, and returned, then your child will not be vaccinated at school)</p> <p>Signature of Parent/Legal Guardian _____</p> <p>Date: month ____ day ____ year ____</p>	<p>I DO NOT GIVE CONSENT to Monmouth County Regional Health Commission for my child named at the top of this form to be vaccinated with this vaccine.</p> <p>Signature of Parent/Legal Guardian _____</p> <p>Date: month ____ day ____ year ____</p>
---	--

Section 4: Permission to Release Information

Placeholder for parental consent for release of data from vaccination record.

Section 5: Vaccination Record

FOR ADMINISTRATIVE USE ONLY

Vaccine	Date Dose Administered	Route	Dose Number (1st or 2nd)	Vaccine Manufacturer	Lot Number	Name and Title of Vaccine Administrator
2009 H1N1	/ /	<input type="checkbox"/> IM <input type="checkbox"/> Intranasal				
2009 H1N1	/ /	<input type="checkbox"/> IM <input type="checkbox"/> Intranasal				